

Nomenclatural Status and Some Additions to the Species Listed in the
Publication, *New Species of Mosquitoes in the Fauna of the*
USSR by A. V. Gutsevich and A. M. Dubitskiy (1981)
(Diptera: Culicidae)

Ronald A. Ward ^{1,2}

ABSTRACT. The taxonomic position of a number of species of Culicidae from the USSR is discussed. The following four species have been recorded from the USSR since 1981: *Aedes* (*Edwardsaedes*) *bekkui*, *Ae.* (*Finlaya*) *versicolor*, *Ae.* (*Ochlerotatus*) *cypriones* and *Culiseta* (*Culiseta*) *nipponica*.

INTRODUCTION

Since the publication of "New species of mosquitoes in the fauna of the USSR" by Gutsevich and Dubitskiy (1981), there have been changes in the status of certain species included in their "List of mosquito species of USSR fauna" (l.c., pp. 98-101). In addition, some earlier changes in their taxonomy of several species were not shown by them. This may be related to the lack of pertinent literature at the time the manuscript was submitted, or due to differences in the interpretation of certain species groups or complexes, viz., the *Anopheles maculipennis* complex.

- ¹ Department of Entomology, Walter Reed Army Institute of Research, Washington, D.C. 20307-5100. Research Associate, Department of Entomology, Smithsonian Institution, Washington, D.C.
- ² The views of the author do not purport to reflect the positions of the Department of the Army or the Department of Defense.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1987		2. REPORT TYPE		3. DATES COVERED 00-00-1987 to 00-00-1987	
4. TITLE AND SUBTITLE Nomenclatural Status and Some Additions to the Species Listed in the Publication, New Species of Mosquitoes in the Fauna of the USSR by A. V. Gutsevich and A. M. Dubitskiy (1981) (Diptera: Culicidae)				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Walter Reed Army Institute of Research, Department of Entomology, Washington, DC, 20307				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT see report					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 7	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

COMMENTS ON RECORDED SPECIES

Genus *Anopheles*

The taxonomy and nomenclature of the *Anopheles maculipennis* complex was summarized by White (1978) who utilized morphological characters (including egg features), genetic crossing experiments and polytene chromosome patterns to redefine the various allopatric and sympatric species. This modern treatment which follows Mayr's (1963, 1969) definitions of species and subspecies has been accepted by most mosquito systematists, but was not mentioned by Gutsevich and Dubitskiy (1981). With respect to the *maculipennis* complex, they state (l.c., p.102): "...we consider them as subspecies, as before, because we do not see any basis for changing our viewpoint which earlier was substantiated in brief (Fauna ..., pp. 90-93) (Gutsevich et al. (1971). An exception may be made for *An. sacharovi*, which differs, although slightly from other members of the "*Anopheles maculipennis* complex" in imagoes, larvae and egg structure. The authors place the greatest reliance upon morphological structures for the separation of species and minimize the value of crossing experiments. *Anopheles maculipennis* Meigen is considered by these authors as a single polytypic species with six Palearctic subspecies.

On the basis of White (1978) and earlier studies, the *An. maculipennis* subspecies *maculipennis*, *messeae* Falleroni, *beklemishevi* Stegnii and Kabanova and *atroparvus* Van Thiel from the USSR should be considered separate species.

Anopheles maculipennis subalpinus Hackett and Lewis (of Gutsevich and Dubitskiy 1981:99) was considered as a subspecies of *melanoon* Hackett by Knight and Stone (1977). Later, White (1978) presented evidence that *subalpinus* represents an alternate egg phenotype of *melanoon* with two kinds of eggs intergrading in some localities. Consequently, *maculipennis subalpinus* should be replaced by the name *melanoon* in the USSR list.

Genus *Aedes*
Subgenus *Aedes*

In 1972, Peus reduced *Ae. rossicus* Dolbeskin, Gorickaja and Mitrofanova to a subspecies of *Ae. esoensis* Yamada. This change was not noted by Gutsevich and Dubitskiy (1981) who considered both of these as subspecies of *Ae. cinereus* Meigen. Tanaka et al. (1975) agreed with the earlier arrangement of Peus (1972) in which *cinereus* and *esoensis* were considered as distinct species, with *Ae. rossicus* as a subspecies of *Ae. esoensis*. They had difficulty with the subspecies concept as used by Gutsevich et al. (1971) as the distribution ranges of the "*cinereus*" subspecies *cinereus*, *eroensis* and *rossicus* overlap with each other in vast areas of the USSR.

Forty-four species of the subgenus *Ochlerotatus* were listed by Gutsevich and Dubitskiy (1981: 99-100). One species, *Ae. refiki* Medschid, was transferred to the monotypic subgenus *Rusticoidus* by Shevchenko and Prudkina (1973). Although this subgenus is well recognized (Knight and Stone 1977), no mention was made of the current status of *Ae. refiki*.

Mezenov (1980) considered *Ae. leucomelas* (Meigen) as a synonym of *Ae. implicatus* Vockeroth. This paper was not mentioned by Gutsevich and Dubitskiy (1981), possibly because their manuscript had been submitted prior to the publication of Mezenov (1980). Gutsevich and Dubitskiy (1981:118) do mention that there are similarities between *Ae. implicatus* and *Ae. leucomelas* and indicate that further study is needed to differentiate the two species. L. T. Nielsen and C. Dahl (personal communication, 1986) consider that they should be retained as separate species; consequently, the synonymy of Mezenov (1980) is not accepted.

The synonymy of *Ae. stramineus* Dubitskiy with *Ae. albineus* Seguy by Danilov (1979) is disputed by Gutsevich and Dubitskiy (1981:121) for the following reasons: The original description of *Ae. albineus* is very brief and it is difficult to make comparisons with USSR material without comparisons with specimens from the type series or type location. It should be mentioned that *Ae. albineus* had been synonymized earlier with *Ae. caspius* (Pallas) (Edwards 1932). Gutsevich and Dubitskiy (1981) think a synonymy with *Ae. campestris* Dyar and Knab is more plausible. Without comparison to the respective three types, these proposed synonyms should be left open.

Aedeia simonini Gutsevich is considered to be a synonym of *Ae. niphadopsis* Dyar and Knab by Danilov (1981). The latter species is only known from the plains and foothills of the northwestern USA (Utah, Idaho, Nevada, Oregon and California). Although the descriptions of the two species are rather similar, the male genitalia, female tarsal claws and the arrangement and number of the comb scales on larval segment VIII are quite different. Without careful comparison of material from both the USSR and the USA, this synonymy should be held in abeyance.

On page 111 of the *Aedes* species key (Gutsevich and Dubitskiy 1981), *Ae. communis* (de Geer) and *Ae. pionips* Dyar are not differentiated by their male genitalia (couplet 41). Danilov (1984a) indicates that he has found two good characters in the male genitalia: differences in the length and shape of the setae on the basistyle apical lobe (apicodorsal lobe of gonocoxite) and shape of the claspette filament apex.

Genus *Culex*
Subgenus *Culex*

Culex pipiens Linnaeus is considered to contain two subspecies (*p. pipiens* and *p. molestus* Forskål) in the USSR. Gutsevich et al. (1971:370) are fully cognizant of the biological character of *Cx. p. molestus*, however they justify the maintenance of subspecific status on behavioral and ecological differences in autogenous-eurygamous populations. This does not take into account the modern concept of species developed by Ernst Mayr and associates over the past 40 years (Mayr 1963). It is advisable to consider *Cx. pipiens molestus* as a behavioral/physiological variant of *Cx. pipiens* without nomenclatorial status (Harbach et al. 1984).

Subgenus

Although *Culex vorax* (Edwards) was placed in synonymy with *Cx. halifaxii* Theobald by Bram (1967), this was not mentioned by Gutsevich et al. (1971) or Gutsevich and Dubitskiy (1981). In the latter paper, reference is only made to the earlier mosquito catalog (Stone et al. 1959) rather than the later edition (Knight and Stone 1977.) The synonymy of Bram (1967) is still considered valid.

Subgenus *Neoculex*

Culex hortensis Ficalbi was transferred from the subgenus *Neoculex* to the subgenus *Maillotia* by Mattingly in 1955. Even though *Cx. hortensis* has been listed under the subgenus *Maillotia* in several mosquito catalogs since 1959, Gutsevich and Dubitskiy (1981) incorrectly retained it in *Neoculex*.

a/ *Culex hayashii* Yamada was transferred from *Neoculex* by Sirivanakarn (1971) to the subgenus *Eumelxomyia* during his reclassification of *Neoculex*. This subgeneric change was not noted.

Genus *Culiseta* Subgenus *Culiseta*

Culiseta subochrea (Edwards) was elevated to specific status by Ribeiro et al. (1977) from a subspecies of *Cs. annulata* (Schrank). This change was also not recorded, possibly due to inaccessible literature.

Subgenus *Culicella*

Although *Culiseta setivalva* (Maslov) was assigned to the subgenus *Culicella* (Maslov 1967, Gutsevich et al. 1971), it was listed by Gutsevich and Dubitskiy (1981) in the subgenus *Culiseta*, possibly through an oversight. Danilov (1984b) reduced *Cs. setivalva* to a synonym of *Cs. fumipennis* (Stephens). If this synonymy is verified, then *Cs. setivalva* should be removed from the USSR fauna and *Cs. fumipennis* added.

Genus *Mansonia*

The generic transfer of *Mansonia richiardii* (Ficalbi) and *Ma. buxtoni* (Edwards) to the genus *Coquillettidia* was not noted. However, not all taxonomists are in agreement with the separation of *Mansonia* into two genera.

ADDITIONS TO THE USSR FAUNA

The following species have been described since the publication of Gutsevich and Dubitskiy (1981), or on the basis of recorded distributions, should be added to the fauna.

1. *Aedes (Edwardsaedes) bekkui* Mogi - Recorded from the south Primorye (Danilov 1985).
2. *Aedes (Finlaya) versicolor* (Barraud) - Recorded from the Talysh foothills, Azerbaijan SSR by Danilov (1978) who also elevated *versicolor* from a variety of *Ae. pulcritarsis* (Rondani) and transferred *versicolor* from the subgenus *Ochlerotatus* to the subgenus *Finlaya*. Although *Ae. versicolor* was previously known from only the adult female collected in Kashmir, the determination of *versicolor* was made by comparing specimens from the USSR with the original description, which lacked illustrations.
3. *Aedes (Ochlerotatus) cypriodes* Danilov and Stupin (1982) - Described from Siberia.
4. *Culiseta (Culiseta) nipponica* LaCasse and Yamaguti - Recorded from the Primorye region (Danilov 1983).

ACKNOWLEDGMENTS

I wish to thank Bruce A. Harrison and E. L. Peyton, Walter Reed Biosystematics Unit, Smithsonian Institution, Washington, D.C. and John F. Reinert, U. S. Army Medical Materiel Development Activity, Fort Detrick, MD, for their constructive comments on this manuscript.

REFERENCES CITED

- Bram, R. A. 1967. Contributions to the mosquito fauna of Southeast Asia. - II. The genus *Culex* in Thailand (Diptera: Culicidae). Contrib. Am. Entomol. Inst. (Ann Arbor) 2(1):1-296.
- Danilov, V. N. 1978. The first finding of *Aedes (Finlaya) versicolor* Barraud (Diptera: Culicidae) in the USSR, description of its female and larva of instar IV. (In Russian) Vestn. Zool. Kiev (3):82-85.
- Danilov, V. N. 1979. Nomenclature, synonymy, differential diagnosis, and distribution of *Aedes (Ochlerotatus) albineus* Seguy (Diptera). (In Russian) Vestn. Zool. Kiev (1):29-35.

- Danilov, V. N. 1981. On the identity of *Aedes (Ochlerotatus) simanini* Gutsevich with *Ae. (O.) niphadopsis* Dyar et Knab (Diptera, Culicidae). (In Russian) Vestn. Zool. Kiev (3):36-40.
- Danilov, V. N. 1983. The species *Culiseta (Culicella) nipponica* LaCasse et Yamaguti new for the fauna of the USSR (Culicidae). (In Russian) Parazitologiya 17:307-309.
- Danilov, V. N. 1984a. The structure of the male genitalia and geographic distribution of the Palearctic mosquitoes of the *Aedes communis* group (Diptera, Culicidae). (In Russian) Vestn. Zool. Kiev (2):38-41.
- Danilov, V. N. 1984b. *Culiseta (Culicella) setivalva* Maslov as a synonym of *C. (C.) fumipennis* Stephens. (In Russian) Parazitologiya 18:313-317.
- Danilov, V. N. 1985. Mosquitoes of the subgenus *Edwardsaedes* of the fauna of the Palearctic with a description of the larva of *Aedes (Edw.) bekkui*. (In Russian) Parazitologiya 19:378-381.
- Danilov, V. N. and V. I. Stupin. 1982. A new species of the mosquito genus *Aedes* Meigen (Diptera, Culicidae) from southern Siberia and Priamuria. (In Russian) Entomol. Obozr. 61:824-829.
- Edwards, F. W. 1932. Genera Insectorum. Diptera. Fam. Culicidae. Fasc. 194, 258pp., Desmet-Verteneuill, Brussels.
- Gutsevich, A. V., A. S. Monchadskii and A. A. Shtakel'berg. 1971. Fauna of the USSR. Diptera. Vol. 3, No. 4. Mosquitoes. Family Culicidae. (In Russian) Akad Nauk SSSR, Zool. Inst. N.S. 100., 384pp.
- Gutsevich, A. V. and A. M. Dubitskiy. 1981. New species of mosquitoes in USSR fauna. (In Russian) Parazitol. Sborn. 30:97-165.
- Harbach, R. E., B. A. Harrison and A. M. Gad. 1984. *Culex (Culex) molestus* Forskål (Diptera: Culicidae): neotype designation, description, variation and taxonomic status. Proc. Entomol. Soc. Wash. 86:521-542.
- Knight, L. K. and A. Stone. 1977. A catalog of the mosquitoes of the world (Diptera: Culicidae). 2nd edition. Thomas Say Found., Entomol. Soc. Am., vol. 6, xi + 611 pp.
- Maslov, A. V. 1967. Blood-sucking mosquitoes of the subtribe Culisetina (Diptera, Culicidae) in world fauna. (In Russian) Akad. Nauk SSSR, Opred. Fauna USSR, No. 93, 182 pp.
- Mattingly, P. F. 1955. Le sous-genre *Neoculex* (Diptera, Culicidae) dans la sous-region mediterraneenne. I-Especie, sous-especie et synonymie nouvelles. Ann. Parasitol. Hum. Comp. 30:374-388.

- Mayr, E. 1963. Animal species and evolution. Harvard Univ. Press, Cambridge, 797 pp.
- Mayr, E. 1969. Principles of systematic zoology. McGraw-Hill, New York, 428pp.
- Mezenev, N. P. 1980. A possible identity of *Aedes leucomelas* Meigen, 1804 and *A. implicatus* Vockeroth, 1954 (Diptera, Culicidae). (In Russian) Parazitologiya 14:206-209.
- Peus, F. 1972. Über das subgenus *Aedes* sensu strictu in Deutschland (Diptera, Culicidae). Z. Angew. Entomol. 72:177-194.
- Ribeiro, H., H. da Cunha Ramos, R. Antunes Capela and C. Alves Pires. 1977. Research on the mosquitoes of Portugal (Diptera, Culicidae) III - Further five new mosquito records. Garcia de Orta, Ser. Zool. 6:51-60, 6 pl.
- Shevchenko, A. K. and N. S. Prudkina. 1973. Morphology of genitals in mosquito males. Report I. Study of genitals of male mosquitoes of the genus *Aedes* (In Russian) Vestn. Zool. (6):40-47.
- Sirivanakarn, S. 1971. Contributions to the mosquito fauna of Southeast Asia. XI. A proposed reclassification of *Neoculex* Dyar based principally on the male terminalia. Contrib. Am. Entomol. Inst. (Ann Arbor) 7(3):62-85.
- Stone, A., K. L. Knight and H. Starcke. 1959. A synoptic catalog of the mosquitoes of the world (Diptera, Culicidae). Thomas Say Found., Entomol.
- Tanaka, K., K. Mizusawa and E. S. Saugstad. 1975. A new species of the genus *Aedes* (*Aedes*) from Japan with synonymical notes on Japanese species of the subgenus *Aedes* (Diptera, Culicidae). Mosq. Syst. 7:41-58, 174-177.
- White, G. V. 1978. Systematic reappraisal of the *Anopheles maculipennis* complex. Mosq. Syst. 10:13-44.